

WIPES

ONE STEP DISINFECTANT CLEANER & DEODORIZER

For Veterinary Use

Reference Sheet

KEEP OUT OF REACH OF CHILDREN

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
viroxanimalhealth.com



EPA Est. No. 74559-CAN-1 EPA Reg. No. 74559-10

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces

It cleans quickly by removing dirt, grime, blood, urine, mold stains, laboratory stains, and other common soils found in animal housing facilities, grooming facilities, kennels, pet stores, veterinary clinics, or other small animal facilities.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile and stainless steel associated with high touch environmental surfaces, animal equipment, animal crates, cages, carts, examination tables, grooming equipment, reception areas, veterinary care surfaces or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

BACTERICIDAL IN 1 MINUTE:

Acinetobacter baumannii (Multi-Drug Resistant)

Bordetella bronchiseptica

Bordetella pertussis

Burkholderia cepacia

Enterobacter aerogenes

Enterobacter cloacae (New Delhi metallo-ß-lactamase 1 (NDM -1))

Enterococcus faecalis (Vancomycin Resistant (VRE))

Enterococcus faecium

Escherichia coli

Escherichia coli O157:H7

Escherichia coli (Extended Spectrum Beta-lactamase (ESBL))

Escherichia coli (New Delhi metallo-ß-lactamase 1 (NDM-1))

Klebsiella pneumoniae

Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))

Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))

Klebsiella pneumoniae (Carbapenem Resistant (CRE))

Legionella pneumophila

Listeria monocytogenes

Pasteurella multocida

Pseudomonas aeruginosa

Pseudomonas aeruginosa (Metallo-Beta lactamase (MBL) positive)

Salmonella enterica (formerly known as Salmonella choleraesuis)

Serratia marcescens

Shigella dysenteriae

Staphylococcus aureus

Staphylococcus aureus Community Acquired Methicillin Resistant Genotype 400 (CA-MRSA 400)

Staphylococcus aureus (Methicillin Resistant (MRSA))

Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))

Staphylococcus aureus (Vancomycin Resistant (VRSA))

Staphylococcus epidermidis (Methicillin Resistant (MRSE))

Staphylococcus pseudintermedius

Stenotrophomonas (Pseudomonas) maltophilia

Streptococcus pneumoniae (Penicillin Resistant (PRSP))

Streptococcus pyogenes

***VIRUCIDAL IN 30 SECONDS:**

Adenovirus Type 8

Avian Influenza A Virus (H7N9)

Avian Influenza A Virus (H3N2)

Cytomegalovirus

Feline Calicivirus

Hepatitis B Virus (Duck Hepatitis B virus as the surrogate)

Hepatitis C Virus (Bovine Viral Diarrhea virus as the surrogate)

Herpes Simplex Virus Type 1

Herpes Simplex Virus Type 2

HIV Type 1 Strain HTLV III_B

Human Coronavirus

Influenza A Virus (Hong Kong Strain)

Influenza B Virus

Norovirus (Feline Calicivirus as the surrogate)

Pandemic Influenza A Virus H1N1

Parainfluenza Virus Type 3

Respiratory Syncytial Virus (RSV)

Rhinovirus Type-37

Rotavirus (WA strain)

Vaccinia Virus

ANIMAL PREMISES *VIRUCIDAL ACTIVITY IN 1 MINUTE, UNLESS OTHERWISE NOTED

Bovine Viral Diarrhea Virus - 30 seconds

Duck Hepatitis B virus - 30 seconds

Equine Influenza A Virus (H3N8) (EIV) - 30 seconds

Feline Calicivirus - 30 seconds

Feline Herpes Virus (Feline Rhinotracheitis)

Feline Infectious Peritonitis Virus (FIPV)

Minute Virus of Mice (Murine Parvovirus)

Mouse Hepatitis Virus

Murine Norovirus - 30 seconds

This product has demonstrated effectiveness against Influenza Type A_2 (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

FUNGICIDAL IN 30 SECONDS, UNLESS OTHERWISE NOTED:

Candida albicans – 1 minute Microsporum canis – 1 minute Trichophyton mentagrophytes

MALODORS

Malodors – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

MOLD/MILDEW

Mildewcidal Activity – kills the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity – controls and prevents the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

TUBERCULOCIDAL IN 1 MINUTE:

Mycobacterium bovis (BCG)

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER IN 10 SECONDS AGAINST:

Acinetobacter baumannii

Campylobacter jejuni

Enterobacter aerogenes

Enterococcus faecalis (Vancomycin Resistant (VRE))

Escherichia coli O157:H7

Listeria monocytogenes

Pseudomonas aeruginosa

Salmonella enterica (formerly known as Salmonella choleraesuis)

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is \underline{not} a baby wipe. Do not flush in toilet.

*KILLS HIV AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY

FLUIDS in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/ body fluids, and in which the surfaces/objects likely to be soiled with blood/ body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV and all other organisms listed on the label.

Disposal of Infectious Material: Cleaning materials used that may contain blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

For Use as a One-Step Cleaner/Disinfectant:

- 1. Pre-clean heavily soiled areas.
- Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- 3. Allow surface to remain visibly wet for 1 minute.
- 4. Wipe surface dry or allow to air dry.

To Kill Fungi:

- 1. Pre-clean heavily soiled areas.
- Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- **3.** All surfaces must remain visibly wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
- 4. Wipe surfaces dry or allow to air dry.

For Use as a Tuberculocide:

- 1. Pre-clean heavily soiled areas.
- 2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- 3. Allow the surface to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
- 4. Wipe surface dry or allow to air dry.
- 5. Change towelette after each use.

To Kill Trichophyton mentagrophytes:

- 1. Pre-clean heavily soiled areas.
- 2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- **3.** All surfaces must remain visibly wet for 30 seconds. Use additional towelettes if needed to ensure the correct wet contact time.
- 4. Wipe surfaces dry or allow to air dry.

To Kill Mold and Mildew:

Pre-clean surfaces. Wipe product onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 4 minutes. Allow to air dry.

To Control Mold and Mildew:

Wipe product onto non-porous surfaces until completely wet. Allow to air dry. Repeat application weekly or when growth reappears.

For use as a Broad Spectrum Non-Food Contact Sanitizer:

- 1. Pre-clean heavily soiled areas.
- 2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- 3. Allow the surface to remain wet for 10 seconds.
- 4. Wipe surface dry or allow to air dry.
- 5. Change towelette after each use

For Use as a Cleaner and/or Deodorizer

Pull towelette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

Animal Housing Facilities in Veterinary Clinics and Hospitals

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, catteries, etc.

- Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
- 2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
- $\textbf{3.} \ \, \textbf{Empty all troughs, racks, and other feeding and watering appliances.}$
- 4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
- 5. All surfaces must remain visibly wet for 1 minute.
- 6. Allow to air dry.
- 7. Ventilate building and other closed spaces.
- 8. Do not house animal or employ equipment until treatment has been absorbed, set or dried.
- Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.